

MARYLAND DEPARTMENT OF HEALTH
PREVENTION AND HEALTH PROMOTION ADMINISTRATION

**MARYLAND MATERNAL
MORTALITY REVIEW
2017 ANNUAL REPORT**

Health – General Article § 13-1207

Larry Hogan
Governor

Boyd K. Rutherford
Lieutenant Governor

Robert R. Neall
Secretary

(This page intentionally left blank)

TABLE OF CONTENTS

Acknowledgements.....1

Background.....2

Methodology.....4

Case Findings.....6

 Cases by Cause of Death Category.....6

 Cases by Timing of Death in Relation to Pregnancy.....7

 Cases by Outcome of Pregnancy8

 Cases by Maternal Race and Ethnicity9

 Cases by Maternal Age9

 Cases by Timing of Prenatal Care Initiation.....10

 Cases by Jurisdiction of Residence and Occurrence.....11

 Preventability of Deaths.....13

Focus on Substance Use Disorder and Overdose Deaths13

2017 MMR Recommendations17

Summary.....19

Appendix: Maryland Maternal Mortality Review Case Discussion Guide20

(This page intentionally left blank)

ACKNOWLEDGEMENTS

This review of maternal deaths would not be possible without the data, cooperation, and expertise of the Maryland Department of Health's Vital Statistics Administration and the Office of the Chief Medical Examiner. The Maternal Mortality Review Program would like to thank the volunteer Committee members for the hours spent in discussion and the serious attention given to this important public health project. The Program is also grateful for the diligent work of the case abstractors in their careful and thorough abstraction of the cases. Special thanks to all those who participated in this year's case review and policy meetings:

<u>Name</u>	<u>Hospital / Affiliation</u>
Pablo Argeles, MD	University of Maryland Baltimore Washington Medical Center
Pedro Arrabal, MD	Sinai Hospital
Robert Atlas, MD	Mercy Medical Center, UMMS
Shayna Banfield, MS, CHES	MedChi, MMR Program Coordinator
Michele Beaulieu, MSW, LCSW-C	University of Maryland School of Social Work
Shobana Bharadwaj, MD	University of Maryland Medical System
Ann Burke, MD	Holy Cross Hospital
Aisha Chaudhry, MS	MedChi, Project Assistant
Diana Cheng, MD	MMR Abstractor
Andreea Creanga, MD, PhD	Johns Hopkins Bloomberg School of Public Health
Sarah Crimmons, MD	University of Maryland Medical System
Stephen Contag, MD	University of Maryland Medical System
Deborah Doerfer, CNM	Johns Hopkins Hospital
Jill Edwardson, MD	Johns Hopkins Bayview Medical Center, MMR Abstractor
Jen Fahey, CNM	University of Maryland Medical System
Casey Fay, MS, CHES	Planned Parenthood of MD
Gia Firth, CNM	Chase Brexton Health Services
Stacy Fisher, MD	University of Maryland Medical System
Lorraine Goldstein, CNM	MMR Abstractor
Katherine Goetzinger, MD	University of Maryland Medical System
Elizabeth Greely, MD	Anne Arundel Medical Center
Asrar Green, RN	Medstar Southern Maryland Hospital
Maureen Grundy, MD	Johns Hopkins Hospital
Mary Carol Jennings, MD, MPH	Johns Hopkins Bloomberg School of Public Health
Clark Johnson, MD	Johns Hopkins Hospital, MMR Committee Chair
Jan Kriebs, CNM	MMR Abstractor
Lorraine Milio, MD	Johns Hopkins Bayview Medical Center, Center for Addiction and Pregnancy, MMR Abstractor
Joseph Morris, MD	Anne Arundel Medical Center
Claudia Richardson, MD	Planned Parenthood of Maryland
Judith Rossiter, MD	University of Maryland St. Joseph Medical Center
Jeanne Sheffield, MD	Johns Hopkins Hospital
Sam Smith, MD	Medstar Franklin Square Medical Center; MedStar Harbor Hospital
Henry Sobel, MD, MBA	Anne Arundel Medical Center
S. Lee Woods, MD, PhD	Maryland Department of Health, MMR Program Director

BACKGROUND

Md. Ann. Code Health-General Art., §13-1203—1207, establishes the Maternal Mortality Review Program (the Program) in the Maryland Department of Health and describes its scope. The purpose of the Program is to: (1) identify maternal death cases; (2) review medical records and other relevant data; (3) determine preventability of death; (4) develop recommendations for the prevention of maternal deaths; and (5) disseminate findings and recommendations to policymakers, health care providers, health care facilities, and the general public.

The Maryland Department of Health (the Department) conducts maternal mortality reviews in collaboration with MedChi, the Maryland State Medical Society. The Department provides funding to MedChi to assist in the maternal mortality review process by obtaining medical records, abstracting cases, and hosting a committee of clinical experts from across the State, the Maternal Mortality Review Committee (the MMR Committee). The MMR Committee provides an in-depth review of maternal deaths to determine pregnancy-relatedness and preventability. The MMR Committee then develops recommendations for the prevention of maternal deaths, and disseminates their findings and recommendations to policy makers, health care providers, health care facilities, and the general public.

Key Definitions

- A **maternal death** is defined by the World Health Organization’s (WHO’s) International Classification of Diseases Ninth and Tenth Revisions (ICD-9 and ICD-10) as “the death of a woman while pregnant or within 42 days of termination of pregnancy, irrespective of the duration and site of the pregnancy, from any cause related to or aggravated by pregnancy or its management but not from accidental or incidental causes.”
- The **maternal mortality ratio or rate (MMR)** is the number of maternal deaths per 100,000 live births.
- A **pregnancy-associated death** is defined by the Centers for Disease Control and Prevention (CDC) as “the death of a woman while pregnant or within one year or 365 days of pregnancy conclusion, irrespective of the duration and site of the pregnancy, regardless of the cause of death.”
- The **pregnancy-associated mortality rate** is the number of pregnancy-associated deaths per 100,000 live births.
- A **pregnancy-related death** is defined by the CDC as “the death of a woman while pregnant or within one year of conclusion of pregnancy, irrespective of the duration and site of the pregnancy, from any cause related to or aggravated by her pregnancy or its management, but not from accidental or incidental causes.”
- The **pregnancy-related mortality rate** is the number of pregnancy-related deaths per 100,000 live births.


The three terms “maternal death,” “pregnancy-associated death,” and “pregnancy-related death,” create a challenge when comparing data from different sources and reports for different jurisdictional entities. The WHO monitors maternal deaths worldwide as a key indicator of population health, and of social and economic development. Maternal deaths are identified solely from information on the death certificate or similar registration of the occurrence and cause of death. Maternal deaths are limited in both the time period and causes considered.

In more developed countries with improved medical care, many deaths related to pregnancy occur beyond 42 days after the end of pregnancy. In 1986, the CDC and the American College of Obstetricians and Gynecologists (ACOG) collaborated to recommend the use of expanded definitions to more accurately identify deaths among women where pregnancy was a contributing factor. This collaboration led to the

definitions for pregnancy-associated and pregnancy-related deaths. Enhanced surveillance methods are necessary to determine pregnancy-associated and pregnancy-related deaths and are discussed below.

Rising Rates of Maternal Mortality

Nationally, maternal mortality has declined dramatically since the 1930s when the MMR was 670 maternal deaths per 100,000 live births. The U.S. MMR was at its lowest level in 1987 at 6.6 maternal deaths per 100,000 live births. However, the MMR has risen since that time, and was 28.7 maternal deaths per 100,000 live births in 2015, the latest year for which national data are available. To compare Maryland's MMR with the national rate, a five-year average is used. This stabilizes the Maryland rate because maternal deaths are relatively infrequent events that may vary considerably year to year, particularly in a small state like Maryland. The Maryland MMR has generally been higher than the national average, although for the period from 2011 to 2015, the Maryland MMR was slightly lower than the national rate (Figure 1). Both are well above the Healthy People 2020 Objective MICH-5 target of 11.4 maternal deaths per 100,000 live births. Between the two 5-year periods shown, the U.S. MMR increased by 37 percent and the Maryland rate increased by 19 percent.



While Maryland's high MMR is concerning, it is also a reflection of the State's intense efforts to more accurately identify maternal deaths since the mid-1990s. To facilitate identification of maternal deaths, the Maryland death certificate was revised in January 2001 to include questions about pregnancy within the year prior to death. This enhanced surveillance resulted in a more than doubling of the number of maternal deaths identified in Maryland compared with data from the 1980s and early 1990s. Whether the actual risk of a woman dying during pregnancy or within 42 days after has increased is unclear. Many studies have shown an increase in chronic health conditions among pregnant women in the United States, including hypertension, diabetes, and heart disease.^{1, 2, 3} These conditions may put pregnant women at higher risk of adverse outcomes.

¹ Kuklina EV, Ayala C, Callaghan WM. Hypertensive disorders and severe obstetric morbidity in the United States: 1998–2006. *Obstet Gynecol.* 2009;113(6):1299–1306.


² Albrecht SS, Kuklina EV, Bansil P et al. Diabetes trends among delivery hospitalizations in the United States, 1994–2004. *Diabetes Care.* 2010;33(4):768–773.

³ Kuklina EV, Callaghan WM. Chronic heart disease and severe obstetric morbidity among hospitalizations for pregnancy in the USA: 1995–2006. *Br J Obstet Gynaecol.* 2011;118(3):345–352.

Racial Disparity

In the U.S., Black women have an MMR more than two and a half times greater than that for White women, a disparity that has persisted since the 1940s. In Maryland, there is a similarly large disparity between the rates among Black and White women.

Figure 2 shows the MMR by race in Maryland for seven overlapping 5-year periods over the past decade. Compared to 2005-2009, the 2011-2015 White MMR in Maryland increased 34 percent and the Black MMR increased 27 percent. The 2011-2015 Black MMR is 2.7 times the White MMR.



METHODOLOGY

Case Identification

Cases for review are limited to women who were residents of Maryland at the time of their death. Maryland residents who died in other states are also included in the MMR case reviews.


Maternal deaths are determined by cause of death information on the death certificates alone. The Maryland death certificate was revised in January 2001 to include questions about pregnancy status, pregnancy outcome, and date of delivery for the 12 months preceding death. Maryland is one of 42 states plus the District of Columbia that now include questions on the death certificate specifically designed to improve identification of maternal deaths. The pregnancy checkbox has significantly increased identification of maternal deaths beyond those recognized by cause of death alone.^{4,5}

⁴ Horon IL. Underreporting of maternal deaths on death certificates and the magnitude of the problem of maternal mortality. *Am J Public Health.* 2005; 95:478-82.

⁵ Horon IL, Cheng D. Effectiveness of pregnancy check boxes on death certificates in identifying pregnancy-associated mortality. *Pub Health Reports.* 2011; 126:195-200.

Pregnancy-associated deaths are identified in one of three ways in Maryland. Individual death certificates are the first method of identifying pregnancy-associated deaths through the use of checkbox questions on the death certificate, or because the cause of death is clearly related to pregnancy (e.g. ruptured ectopic pregnancy or postpartum hemorrhage). The second method of determining pregnancy-associated deaths comes from linking death certificates for women aged 10-50 years with birth certificates and fetal death certificates from the 365 days preceding death to identify additional cases that were not found through examination of death certificates alone. Thirdly, cases reported to the Office of the Chief Medical Examiner are subject to a manual review process to identify evidence of pregnancy in deceased women.

All deaths occurring during pregnancy or within 365 days of pregnancy conclusion are designated as pregnancy-associated and investigated further. Using these three methods, 41 pregnancy-associated deaths were identified in 2015. These cases are reviewed in detail in this report. Figure 3 shows the numbers of pregnancy-associated deaths in Maryland from 2006 to 2015. An average of 40 pregnancy-associated deaths occurred per year during this period.



Case Review

Pregnancy-associated deaths undergo several stages of review. Once cases are identified, medical records are obtained from the hospitals of death and delivery, when applicable. Physician and nurse-midwife abstractors review death certificates, hospital records, and Medical Examiner records for all cases and prepare case summaries that go to the MMR Committee for review. All 2015 pregnancy-associated deaths from all causes (medical, injury, substance use, homicide, and suicide) were reviewed for cause of death, pregnancy-relatedness, and preventability.

Pregnancy-relatedness and potential preventability of the deaths are determined through Committee discussion. The MMR Committee includes obstetric, maternal fetal medicine, nurse-midwifery, nursing, and social work specialties, as well as representatives from the Department's Maternal and Child Health Bureau, Vital Statistics Administration, and the Office of the Chief Medical Examiner. Representatives from all delivery hospitals in Maryland are encouraged to participate. The Committee discussions incorporate the CDC framework for case review outlined in "Strategies to Reduce Pregnancy-Related Deaths: From Identification and Review to Action."⁶ This approach takes into account medical and non-medical factors contributing to maternal death, and examines quality and content of medical care (see Appendix A, *Maryland Maternal Mortality Review Case Discussion Guide*). Cases discussed by the Committee are de-identified and members sign confidentiality agreements. The Committee holds one meeting annually to review issues identified through case reviews and to develop recommendations.


⁶ Berg C, Danel I, Atrash H, Zane S, Bartlett L (Editors). Strategies to reduce pregnancy-related deaths: from identification and review to action. Atlanta: Centers for Disease Control and Prevention; 2001 <<https://stacks.cdc.gov/view/cdc/6537>>.

CASE FINDINGS

A total of 41 pregnancy-associated deaths were identified in 2015 for a pregnancy-associated mortality rate of 55.7 deaths per 100,000 live births. Of the 41 deaths, 12 were determined to be pregnancy-related, while the remaining 29 were either determined not to be related to pregnancy or the relatedness to pregnancy could not be determined. The resulting pregnancy-related mortality rate was 16.3 deaths per 100,000 live births.

Cases by Cause of Death Category



Figure 4 shows the categories of cause of death for pregnancy-associated and pregnancy-related deaths. The leading cause of death among the 41 pregnancy-associated deaths in 2015 was substance use with unintentional overdose, accounting for 12 deaths (29 percent), the greatest number recorded in one year. Injury accounted for five pregnancy-associated deaths (12 percent), suicide for three (seven percent), and homicide an additional one (two percent). The remaining 20 pregnancy-associated deaths (49 percent) were due to natural causes.



Among the 12 pregnancy-related deaths in 2015, the leading cause of death was thrombotic pulmonary embolism, accounting for three deaths (25 percent). An additional seven deaths (58 percent) resulted from other medical causes. The remaining two pregnancy-related deaths (17 percent) resulted from non-natural causes, including one homicide and one suicide.



Cases by Timing of Death in Relation to Pregnancy

Of the 41 pregnancy-associated deaths in 2015, six deaths (15 percent) occurred during pregnancy, ten deaths (24 percent) occurred within 42 days postpartum, and 25 deaths (61 percent) occurred between 43-365 days postpartum (see Figure 5). Among the 12 pregnancy-related deaths, two (17 percent) occurred during pregnancy, eight (66 percent) within 42 days postpartum, and two (17 percent) between 43-365 days postpartum (see Figure 6).




Cases by Outcome of Pregnancy

Among 41 pregnancy-associated deaths in 2015, 33 cases (80 percent) had a live birth, six (15 percent) had co-occurring maternal and fetal deaths, and two cases (five percent) involved a fetal death prior to the mother's death (see Figure 7). Among 12 pregnancy-related deaths, nine cases (75 percent) had a live birth, two (17 percent) had co-occurring maternal and fetal deaths, and one case (eight percent) involved a fetal death (see Figure 8).



Cases by Maternal Race and Ethnicity


Of the 41 pregnancy-associated deaths during 2015, 19 (46 percent) occurred among non-Hispanic White women, 18 (44 percent) among non-Hispanic Black women, three (seven percent) among Asian women, and one (two percent) to a Hispanic woman. Two (17 percent) of the 12 pregnancy-related deaths were among non-Hispanic White women, seven (58 percent) among non-Hispanic Black women, two (17 percent) among Asian women, and one (8 percent) to a Hispanic woman. Pregnancy-associated and pregnancy-related mortality rates by race in 2015 are shown in Figure 9. Rates for racial or ethnic groups with fewer than five deaths are not displayed.




Among 2015 deaths, the pregnancy-associated mortality rate for non-Hispanic Black women was 1.7 times higher than the rate for non-Hispanic White women. Sixty-seven percent of the unintentional overdose deaths were among non-Hispanic White women. Of deaths due to all medical causes, 32 percent occurred among non-Hispanic White women, 10 percent among Asian women, and 58 percent among non-Hispanic Black women.

Cases by Maternal Age


The distribution of pregnancy-associated deaths by maternal age group is shown in Figure 10. The distribution of pregnancy-related deaths by maternal age group is shown in Figure 11.






Cases by Timing of Prenatal Care Initiation

The distributions of pregnancy-associated and pregnancy-related deaths by the trimester when women initiated prenatal care (PNC) are shown in Figures 12 and 13, respectively. Nine (22 percent) of the 41 pregnancy-associated deaths were among women who initiated care in the first trimester of pregnancy. In 15 (37 percent) of the pregnancy-associated deaths, timing of prenatal care was unknown. Among the 12 pregnancy-related deaths, two (17 percent) were among women who received first trimester prenatal care. Timing of prenatal care was unknown in four pregnancy-related deaths (33 percent).





Cases by Jurisdiction of Residence and Occurrence

Thirteen (32 percent) of the 41 pregnancy-associated deaths were among residents of Baltimore City and eight (20 percent) were among residents of Baltimore County (see Figure 14). Sixteen (39 percent) of the deaths occurred in Baltimore City and seven (17 percent) occurred in Baltimore County (see Figure 15).






Figure 15. Number of Pregnancy-Associated Deaths* by Jurisdiction of Occurrence, Maryland, 2015




*Deaths of women from any cause while pregnant or within 365 days of pregnancy
 Data Source: Maryland Department of Health, Vital Statistics Administration

In 2015, three (25 percent) of the 12 pregnancy-related deaths were among residents of Baltimore County, and two (17 percent) each were among residents of Baltimore City and Prince George's County (see Figure 16). Three (25 percent) of the pregnancy-related deaths occurred in Baltimore City and three (25 percent) in Baltimore County (see Figure 17).

Figure 16. Number of Pregnancy-Related Deaths* by Jurisdiction of Residence, Maryland, 2015



*Deaths of women from any cause while pregnant or within 365 days of pregnancy from any cause related to or aggravated by pregnancy
 Data Source: Maryland Department of Health, Vital Statistics Administration



Preventability of Deaths

Of the 41 pregnancy-associated deaths, 33 (80 percent) were determined to be preventable or potentially preventable. In two cases (5 percent), preventability could not be determined, and six cases (15 percent) were considered unpreventable deaths. Among the 12 pregnancy-related deaths, ten (83 percent) were thought to be preventable or potentially preventable. Two cases (17 percent) were considered unpreventable deaths.

All of the unintentional overdose deaths were considered potentially preventable, as were the three suicide deaths and the one homicide death. Four of the five injury deaths were also considered potentially preventable. The six deaths considered unpreventable involved medical causes of death, including thrombotic pulmonary embolism, cardiovascular conditions, postpartum autoimmune disorders, and cancer.

FOCUS ON SUBSTANCE USE DISORDER AND OVERDOSE DEATHS


In 2015 for the third consecutive year, drug overdose was the leading cause of pregnancy-associated death in Maryland. Twelve of 41 deaths (29 percent) resulted from substance use and unintentional overdose. All of these deaths involved opioids. In 10 of the 12 cases (83 percent) of pregnancy-associated death resulting from substance use and unintentional overdose, two or more drugs were found by postmortem toxicology testing. Three overdose deaths (25 percent) involved the potent opioid fentanyl. A benzodiazepine was present in four cases (33 percent), and alcohol was detected in two cases. The risk of fatal overdose is substantially increased when opioids are combined with other central nervous system depressants such as benzodiazepines or alcohol.

The average age at death among the 12 pregnancy-associated deaths resulting from substance use and unintentional overdose was 32 years (range 23 to 41 years). Eight deaths (67 percent) were among non-Hispanic White women, four (33 percent) among non-Hispanic Black women. Only one woman was pregnant at the time of death. The other 11 women (92 percent) had delivered live born infants, and the average timing of death was 166 days postpartum. Only two deaths occurred in the traditional postpartum period up to 42 days after the conclusion of pregnancy, and nine occurred between 43 and 365 days postpartum. Eleven of the 12 women (92 percent) had a known history of substance use. In all 12 cases,

there was a history of one or more mental health diagnosis, with depression documented in 11 cases, anxiety in ten cases, and bipolar disorder in three cases.

Multiyear Review of Overdose Deaths

To better understand factors involved in overdose deaths, a review of pregnancy-associated deaths in Maryland from 2010 to 2015 was undertaken. Over this six-year period, substance use and unintentional overdose was the leading cause of pregnancy-associated death, accounting for 46 of 228 pregnancy-associated deaths (20 percent). Figure 18 shows the number of unintentional overdose deaths by year, with the highest number of cases occurring in 2015. As shown, very few of these cases were considered pregnancy-related.



Of the 46 overdose deaths, 44 (96 percent) involved opioids (one of the remaining two cases involved alcohol, and the other involved alcohol plus the amphetamine methylone). Table 1 shows the specific opioid(s) identified by toxicology testing at the time of death in the 44 cases involving opioids. The most frequently detected opioid was morphine, a metabolite of heroin, followed by methadone and oxycodone. Fentanyl was detected in three cases from 2015, one case from 2014, and none occurring earlier. In 42 (91 percent) of the 46 overdose deaths, two or more drugs were detected by postmortem testing. In 11 (26 percent) of the 42 multiple drug cases, two or three different opioids were identified. Benzodiazepines were detected in 14 (33 percent) and alcohol in 11 (26 percent) of the cases involving multiple drugs.

**Table 1. Opioid Identified Postmortem,
Pregnancy-Associated Unintentional Overdose Deaths,
Maryland, 2010-2015**

Opioid	Number of cases (n=44)
Morphine (heroin)	16
Methadone	13
Oxycodone	8
Unspecified opioid	5
Fentanyl	4
Tramadol	3
Codeine	2
Buprenorphine	1
Hydrocodone	1
Hydromorphone	1
Meperidine	1
Oxymorphone	1

Data Source: Maryland Department of Health
NOTE: The values in the table do not add up to the sample size of 44 because multiple drugs can be detected in a single case.

The average age at death was 29 years among the 46 unintentional overdose deaths from 2010 to 2015. Thirty-eight (83 percent) of these deaths were among non-Hispanic White women and eight (17 percent) among non-Hispanic Black women. Seven women (15 percent) were pregnant at the time of death and five (11 percent) had a spontaneous abortion or fetal demise prior to death. The remaining 34 women (74 percent) delivered live born infants. Only four deaths (nine percent) occurred at 42 days or less postpartum; seven (15 percent) were pregnant at death; and the remaining 35 (76 percent) occurred between 43 and 365 days postpartum. The average timing of death was 197 days postpartum. In 39 cases (85 percent), one or more mental health diagnosis was documented. Anxiety was diagnosed in 30 cases (65 percent), depression in 29 (63 percent) and bipolar disorder in 16 (35 percent). Forty-two (91 percent) of the women who died of overdose had a known history of substance use and twenty-two (48 percent) had documentation of some substance use treatment.

In Table 2, the 46 overdose cases are compared with the 182 non-overdose cases that occurred between 2010 and 2015. Average age at death was comparable in both groups. However, the racial distribution is strikingly different, with a preponderance of non-Hispanic White women among the overdose deaths and overrepresentation of non-Hispanic Black women among the non-overdose deaths. A similar percentage of women were pregnant at the time of death in both groups, but deaths after the conclusion of pregnancy occurred on average much later among the overdose group. Pregnancy outcome was similar in both groups, with 74 percent of pregnancies among the overdose group and 69 percent among the non-overdose group resulting in a live birth. Timing of prenatal care initiation was similar, with more than half of women in both groups starting prenatal care in the first or second trimester.

There were large differences, however, between the two groups related to several behavioral health factors. Women who died of overdose were more than four-times as likely as women who died of other causes to have a known history of substance use (91 percent vs. 19 percent). Women who died of overdose were more than four-times as likely to have one or more mental health diagnosis (85 percent vs. 19 percent), and were more than three-times as likely to smoke (85 percent vs. 24 percent). Also, 44 of 46 overdose deaths (96 percent) were considered preventable or potentially preventable, compared with 56 percent of the non-overdose deaths.

Table 2: Incident Characteristics of Pregnancy-Associated Deaths, Maryland, 2010-2015		
Characteristic	Overdose Deaths (n=46)	Non-overdose Deaths (n=182)
Data presented as mean ± standard deviation, or number (%)		
Demographics		
Average age at death (years)	29 ±5	31 ±7
White non-Hispanic	38 (83)	68 (37)
Black non-Hispanic	8 (17)	89 (49)
Other non-Hispanic	0 (0)	13 (7)
Hispanic	0 (0)	12 (7)
Timing of death		
Pregnant at death	7 (15)	35 (19)
0-42 days postpartum	4 (9)	74 (41)
43-365 days postpartum	35 (76)	72 (40)
Average days postpartum	197 ±93	103 ±117
Pregnancy outcome		
Live born infant	34 (74)	125 (69)
Co-occurring maternal-fetal deaths	7 (15)	35 (19)
Spontaneous abortion / fetal death	5 (11)	14 (8)
Prenatal care initiation		
1 st trimester	16 (35)	76 (42)
2 nd trimester	13 (28)	18 (10)
3 rd trimester	2 (4)	4 (2)
No prenatal care	5 (11)	14 (8)
Termination or very early pregnancy	1 (2)	6 (3)
Unknown	9 (20)	64 (35)
Behavioral health / social factors		
Known history of substance use	42 (91)	34 (19)
Any history of substance use treatment	22 (48)	15 (8)
Smoking	39 (85)	43 (24)
Mental health diagnosis(es)	39 (85)	35 (19)
Intimate partner violence	5 (11)	17 (9)
Preventability		
Death preventable / potentially preventable	44 (96)	101 (55)

Data Source: Maryland Department of Health, Vital Statistics Administration

2017 MMR RECOMMENDATIONS

Unintentional overdose deaths and other behavioral health issues were reviewed in the 2015 MMR Report and detailed recommendations were put forward to address these issues. These recommendations continue to be relevant, as unintentional overdose remains the leading cause of pregnancy-associated death in Maryland. The Committee, therefore, maintains its support of the recommendations related to substance use disorder and unintentional overdose made in the 2015 MMR Report. These recommendations are presented below and the full 2015 report is available at: http://phpa.dhmh.maryland.gov/mch/Documents/2015MMR_FINAL.pdf.

2015 MMR Recommendations - Overdose Deaths	Action Items
<ul style="list-style-type: none"> • Promote universal screening at least once during pregnancy, at delivery, and postpartum for substance use, mental health, and intimate partner violence conditions. • Document screening tools used, referrals given, and treatment plans in perinatal records. • Reduce unintended pregnancy and encourage reproductive life planning. • Improve communication and collaboration between providers of prenatal care and other providers (mental health, substance use, domestic violence, primary care, oral health, etc.). • Promote interdisciplinary case management among substance use, mental health, and intimate partner violence programs. • Improve safe opioid prescribing practices. • Encourage Prescription Drug Monitoring Program (PDMP) utilization by providers. • Encourage naloxone co-prescribing and 3rd party prescribing. • Inform substance use treatment providers about perinatal health. 	<ul style="list-style-type: none"> • Create and disseminate a resource list of valid screening tools for substance use, mental health, and intimate partner violence. • Create and disseminate a resource list of referral service options by Maryland jurisdiction. • Strive for a single point of contact for behavioral health services to facilitate providers' accessing referral sources. • Promote integration of reproductive life planning and preconception counseling into health care visits by all disciplines. • Encourage use of Long Acting Reversible Contraception for women who indicate they do not desire to become pregnant. • Promote the importance of establishing linkages and relationships to ongoing care during the perinatal and postpartum period. • Facilitate obtaining medical records from behavioral health service providers so that the obstetric chart has comprehensive information of the patient's behavioral health care. • Establish a liaison between the MMR Program and the Heroin and Opioid Emergency Task Force, the Task Force on Maternal Mental Health, and intimate partner violence programs. • Raise provider awareness about substance use during pregnancy and promote current resources and trainings. • Educate providers on the use and importance of the PDMP. • Train providers, patients, and families on naloxone use and response to opioid overdose. • Develop a consultation resource about perinatal and reproductive health issues for substance use treatment providers.

In addition, the Committee would like to develop a more consistent and identifiable means of disseminating maternal mortality review findings and recommendations to the provider community. The Committee will develop a standardized **Provider Alert** that can be used to disseminate information about specific causes of maternal deaths in Maryland, and recommendations to improve practices and prevent future deaths. Based on the current review of maternal deaths, the MMR Committee puts forward the following additional recommendations:

Substance Use Disorders and Unintentional Overdose

Recommendation 1:

Increase provider awareness of the contribution of substance use and unintentional overdose to maternal mortality in Maryland.

Action: Develop a *Provider Alert* on substance use as a key contributor to maternal mortality in the State and distribute it widely to obstetric care providers, substance use treatment providers, delivery hospital leadership and staff, and professional organizations.

Recommendation 2:

Support the Maryland Department of Health efforts to coordinate the care of pregnant women with substance use disorders.

Action: Assist with on-going work of the Substance Abuse and Mental Health Services Administration-sponsored Maryland Coalition for the Safe Care of Substance Exposed Infants and Their Families to examine current resources and develop a State plan for care of pregnant and postpartum women with substance use disorders and a safe plan of care for their infants.

Action: Establish collaboration with the Maryland Department of Health's Behavioral Health Administration to compile a list of substance use treatment providers by jurisdiction and distribute it widely to women's health care providers, obstetric care providers, and delivery hospitals.

Late Postpartum Deaths - Care Transitions Beyond 42 Days Postpartum

Recommendation:

Increase provider awareness that: 1) the majority of pregnancy-associated deaths, including those from unintentional overdose, occur well beyond 42 days postpartum; 2) continuity of maternal medical and behavioral health care beyond the traditional six-week postpartum period is critical in preventing late maternal deaths; and 3) the transitional period from perinatal care to primary and/or specialty care is a time of particular health vulnerability for women.

Action: Develop a *Provider Alert* on the importance of postpartum care transition, encouraging that every woman have a documented postpartum care plan and care team identified during the prenatal period.

Action: Make available to prenatal care providers and delivery hospitals resources from and linkage to the Council on Patient Safety in Women's Healthcare's "*Postpartum Care Basics for Maternal Safety*."

Seat Belt Use in Motor Vehicle Collisions

Recommendation:

Encourage increased seatbelt use during pregnancy and awareness by obstetric providers of the importance of seatbelt use during pregnancy.

Action: Develop a *Provider Alert* on seat belt use as key factor in preventing maternal deaths in motor vehicle collisions.

Action: Disseminate ACOG materials, including patient education materials, on the correct use of seatbelts during pregnancy to obstetric care providers and delivery hospitals.

SUMMARY

Maryland's maternal mortality rate is now just below the national average, but remains substantially higher than the Healthy People 2020 goal of 11.4 deaths per 100,000 live births. This in part reflects efforts in the State to accurately identify maternal deaths. Enhanced surveillance methods include questions on the death certificate about pregnancy within the year prior to death, linkage of women's death certificates with birth and fetal death certificates from the previous year, review of medical examiner records, and detailed case review by the MMR Committee.

Forty-one pregnancy-associated deaths were identified in 2015. Twelve (29 percent) of these cases were determined to be pregnancy-related, with the cause of death related to or aggravated by the pregnancy or its management. The leading cause of pregnancy-associated death for the third consecutive year was substance use and unintentional overdose. Thrombotic pulmonary embolism was the leading cause of pregnancy-related death. A majority of these deaths (80 percent of pregnancy-associated deaths and 83 percent of pregnancy-related deaths) were considered preventable or potentially preventable.

In this report, the MMR Committee focused its recommendations on unintentional overdose deaths and improving its dissemination of maternal mortality review findings and recommendations to the provider community. The MMR Committee will continue to promote communication and collaboration among all providers caring for pregnant and postpartum women in an effort to reduce pregnancy-associated deaths in Maryland.

Appendix

Maryland Maternal Mortality Review Case Discussion Guide

Date: _____ Case # _____

Purpose: To review pregnancy-associated deaths in order to classify cases, identify trends in mortality, and develop recommendations for systems change.

Case Definition: Death of a woman while pregnant or within 365 days of pregnancy conclusion

1. Medical Care and Non-medical Causes Underlying the Death

Quality/content of medical care

- Preventive services
- Community and patient education
- Nutrition, substance use, and social services
- Preconception services
- Prenatal care
- Labor and delivery services
- Postpartum care and follow-up
- Management & treatment
- Diagnostic procedures
- Medical interventions
- Patient education and follow-up

Non-medical (social) causes underlying the death

- Intendedness of pregnancy
- Woman's and her family's knowledge about pregnancy & its possible complications
- Timeliness on the part of the woman in recognizing a problem & taking action
- Accessibility/acceptability of healthcare (cultural/experience/financial/geographic/transportation/logistic)
- Cultural competence and communication skills of health care providers
- Woman's adherence or non-adherence to medical advice and health interventions

2. Issues specific to this case

Individual Behavior: _____

Provider Practice: _____

Institutional/ Systems Issues: _____

Additional issues: _____

Sources of Information: _____

Information Missing: _____

1. Type of Case:

Pregnancy-related (causes related to or aggravated by pregnancy or its management)

Not Pregnancy-related (cause unrelated to pregnancy)

Undetermined

Due to: _____

2. This case was:

Preventable (individual provider institutional/systems issues)

Potentially Preventable (individual provider institutional/systems issues)

Undetermined

Not Preventable

3. Resources or services needed but not used or not available:

4. Recommendation(s) to address issues in this case:

